

Critical phenomena of the hard-sphere lattice gas on the simple cubic lattice

Atsushi Yamagata

Department of Physics, Tokyo Institute of Technology, Oh-okayama, Meguro-ku, Tokyo 152, Japan

and

Department of Physics, Tokyo Metropolitan University, Minami-ohsawa 1-1, Hachioji-shi, Tokyo 192-03, Japan

Running title Critical phenomena of hard-sphere lattice gas

Keywords Hard-sphere lattice gas, Critical phenomena, Monte Carlo method

PACS classification codes 02.70.Lq, 64.60.Cn, 68.35.Rh

Abstract

We study the critical phenomena of the hard-sphere lattice gas on the simple cubic lattice with nearest neighbour exclusion by the Monte Carlo method. We get the critical exponents, $\beta/\nu = 0.313(9)$ and $\gamma/\nu = 2.37(2)$, where β is the critical exponent for the staggered density, γ for the staggered compressibility, and ν for the correlation length.

1 Introduction

Atoms of a hard-sphere lattice gas occupy sites of a lattice and interact with infinite repulsion of nearest neighbour pairs. The grand partition function is

$$\Xi_V(z) = \sum_N z^N Z_V(N), \quad (1)$$

where z is an activity and $Z_V(N)$ is the number of configurations in which there are N atoms in the lattice of V sites. At $z = +\infty$ a ground state configuration is that the atoms occupy all the sites of one sublattice and the other is vacant. There is no atom at $z = 0$. A continuous phase transition occurs at a critical activity. There are many studies of the system by various methods: series expansions [1, 2, 3, 4, 5], finite-size scaling and transfer matrix [6], Bethe and ring approximations [7], transfer matrix [8, 9, 10], corner transfer matrix and series expansions [11], exact calculations [12], and Monte Carlo simulations [13, 14].

The critical activity is obtained on various lattice. The square lattice: 3.80(2) [4], 3.80(4) [9], 3.7966(3) [10], 3.7962(1) [11], and 3.796255174(3) [15]. The triangular lattice: 11.12(11) [9], 11.05(15) [5], and $\frac{1}{2}(11 + 5\sqrt{5}) = 11.090\dots$ [12]. The simple cubic lattice: 1.09(7) [5] and 1.0588(3) [14]. The body-centred cubic lattice: 0.77(5) [5] and 0.73(2) [16, 17]. The critical exponents on the square lattice are as follows. The thermal exponent, y_t , is 1.000000(1), the magnetic exponent, y_h , is 1.875000(1) [6], $\beta/\nu = 0.125(5)$, and $\gamma/\nu = 1.74(2)$ [13] where β , γ , and ν are the critical exponents for the order parameter, the staggered compressibility, and the correlation length, respectively. There are exact values on the triangular lattice [12, 18]: $\alpha = 1/3$ for the compressibility, $\beta = 1/9$, $\mu = 5/6$ for the interfacial tension, and $\nu = 5/6$. The value, 0.61071(2) [19], of the fourth-order cumulant [20] of the order parameter on the square lattice agrees with those of the Ising model, 0.611(1) [21], 0.6103(7) [22], 0.611(1), and 0.610690(1) [19]. These results support that the hard-sphere lattice gas on the square lattice belongs to the universality class of the two-dimensional Ising model and that on the triangular lattice does to one of the three-state Potts model in two dimensions [23]. There is no result of critical exponents in three-dimensional hard-sphere lattice gas.

We carry out Monte Carlo simulations of the hard-sphere lattice gas on the simple cubic lattice. Using the finite-size scaling, we estimate the critical

exponents. In the next section we define physical quantities measured. In section 3 we describe procedure of analyses and present Monte Carlo results. A summary is given in section 4.

2 Monte Carlo simulations

We use the Metropolis Monte Carlo technique [24, 25] to simulate the hard-sphere lattice gas (1) on the simple cubic lattice of V sites, where $V = L \times L \times L$ ($L = 2 \times n$, $n = 2, 3, \dots, 30$), under fully periodic boundary conditions. According to Meirovitch [13], we adopt the grand canonical ensemble. The algorithm is described in the references [13, 14].

We start each simulation with a ground state configuration at the critical activity, $z_c = 1.0588$ [14]. The pseudorandom numbers are generated by the Tausworthe method [26, 27]. We measure physical quantities over 1.2×10^6 Monte Carlo steps per site (MCS/site) for $L \geq 44$ and 10^6 MCS/site for $L \leq 42$ after discarding 5×10^4 MCS/site to attain equilibrium. We have checked that simulations from the ground state configuration and no atom one gave consistent results. Each run is divided into M blocks. Let us the average of a physical quantity, O , in each block $\langle O \rangle_i$; $i = 1, 2, \dots, M$. The expectation value is

$$\overline{\langle O \rangle} = \frac{1}{M} \sum_{i=1}^M \langle O \rangle_i.$$

The standard deviation is

$$\Delta \langle O \rangle = \left(\overline{\langle O \rangle^2} - \overline{\langle O \rangle}^2 \right)^{1/2} / \sqrt{M-1}.$$

We have taken $M = 12$ for $L \geq 44$ and $M = 10$ for $L \leq 42$.

Let us define an order parameter by

$$R = 2(N_A - N_B)/V$$

where N_A (N_B) is the number of the atoms in the A (B)-sublattice and $N = N_A + N_B$. We measure the staggered densities,

$$m^\dagger = \overline{\langle |R| \rangle} \tag{2}$$

and

$$m^{\dagger'} = \overline{\langle R^2 \rangle^{1/2}}, \tag{3}$$

the staggered compressibilities,

$$\chi^\dagger = V(\overline{\langle R^2 \rangle} - \langle |R|^2 \rangle)/4 \quad (4)$$

and

$$\chi^{\dagger'} = V\overline{\langle R^2 \rangle}/4, \quad (5)$$

and the fourth-order cumulant of R [20],

$$U = 1 - \frac{1}{3} \frac{\overline{\langle R^4 \rangle}}{\langle R^2 \rangle^2}. \quad (6)$$

3 Monte Carlo results

We estimate a critical exponent and an amplitude by using the finite-size scaling [28, 29, 30]. For a physical quantity, O , we use the nonlinear chi-square fitting [31] with a function of L ,

$$O(L) = A L^p,$$

where p and A are fitting parameters. $p = -\beta/\nu$ when $O = m^\dagger$ or $m^{\dagger'}$. $p = \gamma/\nu$ when $O = \chi^\dagger$ or $\chi^{\dagger'}$. We calculate the chi-square per degrees of freedom, χ^2/DOF , and the goodness of fit, Q , [31] for the data set of the sizes $L = L_{\min}, L_{\min} + 2, \dots, L_{\max} - 2$, and $L_{\max}$. The values of $L_{\min}$ and $L_{\max}$ are selected so that the difference between χ^2/DOF and 1 is the smallest.

We get the results as follows. For $m^\dagger$ defined by (2), $\beta/\nu = 0.28(1)$, $A = 0.39(2)$, $\chi^2/\text{DOF} = 1.21$, and $Q = 0.27$. For $m^{\dagger'}$ defined by (3), $\beta/\nu = 0.313(9)$, $A = 0.48(2)$, $\chi^2/\text{DOF} = 1.22$, and $Q = 0.26$. For $\chi^\dagger$ defined by (4), $\gamma/\nu = 1.96(2)$, $A = 0.040(3)$, $\chi^2/\text{DOF} = 0.98$, and $Q = 0.47$. For $\chi^{\dagger'}$ defined by (5), $\gamma/\nu = 2.37(2)$, $A = 0.058(4)$, $\chi^2/\text{DOF} = 1.21$, and $Q = 0.28$. $L_{\min} = 36$ and $L_{\max} = 60$ for $m^\dagger$, $m^{\dagger'}$, and $\chi^{\dagger'}$. $L_{\min} = 30$ and $L_{\max} = 54$ for $\chi^\dagger$.

The values of the critical exponents of $m^{\dagger'}$ and $\chi^{\dagger'}$ satisfy the scaling relation, $2\beta + \gamma = d\nu$ where d is the dimensions, within errors. Thus we adopt them as the reliable estimates. Figure 1 shows the size dependence of $m^{\dagger'}$ at $z = z_c$. The solid line indicates $0.48L^{-0.313}$. Figure 2 shows the size dependence of $\chi^{\dagger'}$ at $z = z_c$. The solid line indicates $0.058L^{2.37}$. Figure 3 shows the size dependence of U defined by (6) at $z = z_c$. We cannot extract a reliable estimate in $L = +\infty$ since we do not know the correction to scaling of this system. It will have a value between 0.54 and $2/3$.

4 Summary

We carry out the Monte Carlo simulations of the hard-sphere lattice gas on the simple cubic lattice with nearest neighbour exclusion under fully periodic boundary conditions. Using the finite-size scaling, we obtain the critical exponents, $\beta/\nu = 0.313(9)$ and $\gamma/\nu = 2.37(2)$. They satisfy the scaling relation, $2\beta + \gamma = d\nu$.

The critical exponents of the three-dimensional Ising model have been estimated, $\beta/\nu = 0.518(7)$ and $\gamma/\nu = 1.9828(57)$ [32]. These values do not agree with our results of the hard-sphere lattice gas. The value of the fourth-order cumulant of the Ising model on the simple cubic lattice under fully periodic boundary conditions is as follows. 0.44(2) [20], 0.4677(23) [33], and 0.4637(13) [34]. We see that these values are smaller than those of the hard-sphere lattice gas as seen in figure 3. While there is evidence that the hard-sphere lattice gas on the square lattice belongs to the Ising universality class in two dimensions as described in section 1, it does not seem that the hard-sphere lattice gas on the simple cubic lattice falls into the universality class of the three-dimensional Ising model.

Acknowledgements

We have carried out the simulations on the HITAC S-3600 computer under the Institute of Statistical Mathematics Cooperative Research Program (94-ISM-CRP-43 and 95-ISM-CRP-37) and on two personal computers with the 486DX2/66MHz CPU and the Linux operating system (SLS 1.03 + JE 0.9.3, Slackware 2.0.0 + JE 0.9.5). This study was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture, Japan.

References

- [1] C. Domb, *Nuovo Cimento* 9 Suppl. (1958) 9.
- [2] H.N.V. Temperley, *Proc. Phys. Soc.* 74 (1959) 183.
- [3] D.M. Burley, *Proc. Phys. Soc.* 75 (1960) 262.
- [4] D.S. Gaunt and M.E. Fisher, *J. Chem. Phys.* 43 (1965) 2840.
- [5] D.S. Gaunt, *J. Chem. Phys.* 46 (1967) 3237.
- [6] G. Kamieniarz and H.W.J. Blöte, *J. Phys. A* 26 (1993) 6679.
- [7] D.M. Burley, *Proc. Phys. Soc.* 77 (1961) 451.
- [8] L.K. Runnels, *Phys. Rev. Lett.* 15 (1965) 581.
- [9] L.K. Runnels and L.L. Combs, *J. Chem. Phys.* 45 (1966) 2482.
- [10] F.H. Ree and D.A. Chesnut, *J. Chem. Phys.* 45 (1966) 3983.
- [11] R.J. Baxter, I.G. Enting, and S.K. Tsang, *J. Stat. Phys.* 22 (1980) 465.
- [12] R.J. Baxter, *J. Phys. A* 13 (1980) L61.
- [13] H. Meirovitch, *J. Stat. Phys.* 30 (1983) 681.
- [14] A. Yamagata, *Physica A* 215 (1995) 511.
- [15] H.W.J. Blöte and X.-N. Wu, *J. Phys. A* 23 (1990) L627.
- [16] D.P. Landau, *Phys. Rev. B* 16 (1977) 4164.
- [17] Z. Ràcz, *Phys. Rev. B* 21 (1980) 4012.
- [18] R.J. Baxter and P.A. Pearce, *J. Phys. A* 15 (1982) 897.
- [19] G. Kamieniarz and H.W.J. Blöte, *J. Phys. A* 26 (1993) 201.
- [20] K. Binder, *Z. Phys. B* 43 (1981) 119.
- [21] A.D. Bruce, *J. Phys. A* 18 (1985) L873.

- [22] D. Nicolaides and A.D. Bruce, J. Phys. A 21 (1988) 233.
- [23] F.Y. Wu, Rev. Mod. Phys. 54 (1982) 235; 55 (1983) 315 (erratum).
- [24] K. Binder, Monte Carlo Methods in Statistical Physics, K. Binder, ed. (Springer, Berlin, 1979) p. 1.
- [25] K. Binder and D. Stauffer, Applications of the Monte Carlo Method in Statistical Physics, 2nd Ed., K. Binder, ed. (Springer, Berlin, 1987) p. 1.
- [26] N. Ito and Y. Kanada, Supercomputer 5 (1988) 31.
- [27] N. Ito and Y. Kanada, Proceedings of Supercomputing '90 (IEEE Computer Society Press, Los Alamitos, 1990) p. 753.
- [28] M.E. Fisher, Critical Phenomena, Proc. 1970 Enrico Fermi Summer School, Vol. 51, M.S. Green, ed. (Academic Press, New York, 1970) p. 1.
- [29] M.N. Barber, Phase Transitions and Critical Phenomena, Vol. 8, C. Domb and J.L. Lebowitz, eds. (Academic Press, London, 1983) p. 145.
- [30] V. Privman, Finite Size Scaling and Numerical Simulation of Statistical Systems, V. Privman, ed. (World Scientific, Singapore, 1990) p. 1.
- [31] W.H. Press, S.A. Teukolsky, W.T. Vetterling, and B.P. Flannery, Numerical Recipes in C: The Art of Scientific Computing, 2nd Ed. (Cambridge University Press, Cambridge, 1992) Chapters 6 and 15.
- [32] A.M. Ferrenberg and D.P. Landau, Phys. Rev. B 44 (1991) 5081.
- [33] P.-Y. Lai and K.K. Mon, Phys. Rev. B 40 (1989) 11120.
- [34] H.W.J. Blöte and G. Kamieniarz, Physica A 196 (1993) 445.

Figure captions

Figure 1 Size dependence of the staggered density, $m^\dagger$, defined by (3) at $z = z_c$. The solid line indicates $0.48L^{-0.313}$. Errors are less than the symbol size.

Figure 2 Size dependence of the staggered compressibility, $\chi^\dagger$, defined by (5) at $z = z_c$. The solid line indicates $0.058L^{2.37}$. Errors are less than the symbol size.

Figure 3 Size dependence of the fourth-order cumulant, U , defined by (6) at $z = z_c$.




Figure 1
 Atsushi Yamagata




Figure 2

Atsushi Yamagata




Figure 3
 Atsushi Yamagata